

Comment on “Exact bosonization for an interacting Fermi gas in arbitrary dimensions”

D. Galanakis,¹ S. Yang,¹ F.F. Assaad,² M. Jarrell,¹ P. Werner,³ and M. Troyer³

¹*Department of Physics and Astronomy, Baton Rouge, LA, USA*

²*Institut für theoretische Physik und Astrophysik,*

Universität Würzburg, Am Hubland, D-97074, Germany

³*Theoretische Physik, ETH Zurich, 8093, Zurich, Switzerland*

In a recent Letter [1] Efetov *et al.* propose an exact mapping of an interacting fermion system onto a new model that is supposed to allow sign-problem free Monte Carlo simulations. In this Comment, we show that their formalism is equivalent to the standard approach of Blackenbecker, Scalapino and Sugar (BSS) [2] for fermionic systems and has the same sign statistics and minus sign problem.

Our first observation is that the partition function for a given configuration of the auxiliary fields ϕ is the same in the standard formulation Z_f [Eq. (8) in Ref. [1]] and in their new bosonized scheme Z_b [Eq. (9)]:

$$Z_f[\phi] = Z_b[\phi]. \quad (1)$$

This observation is trivial in the limit of the time step $\Delta\tau \rightarrow 0$, where both schemes reproduce the same partition function. Since Z_f can be negative also in this limit [2], Z_b is also not sign-positive. Both Z_f and Z_b are positive if ϕ is a smooth path [2], but restricting the configuration space to smooth paths amounts to a semi-classical approximation.

We next show that $Z_f[\phi] = Z_b[\phi]$ also holds for finite $\Delta\tau$ and piecewise constant paths where the field $\phi_r(\tau)$ is constant on the interval $[(l+1)\Delta, l\Delta]$. Efetov *et al.*'s Eq. (9) is equivalent to $Z_b[\phi] = \text{Tr}(e^{-\beta H_0} U_I(\beta, 0))$ where $\frac{\partial}{\partial \tau_1} U_I(\tau_1, \tau_0) = -H_{I,1}(\tau_1) U_I(\tau_1, \tau_0)$ and $H_{I,1}(\tau) = -\sum_r \phi_r(\tau) e^{\tau H_0} m_{z,r} e^{-\tau H_0}$. Since $U_I(\tau_2, \tau_0) = U_I(\tau_2, \tau_1) U_I(\tau_1, \tau_0)$, $Z_b[\phi] = \text{Tr}[e^{-\beta H_0} \prod_{n=0}^N U_I(l\Delta, (l-1)\Delta)]$. We are left with the task of evaluating $U_I(l\Delta, (l-1)\Delta)$ on the l^{th} time interval where the fields are constant and take the values ϕ_{rl} . For this constant in time field configuration, $U_I(l\Delta, (l-1)\Delta) = e^{l\Delta H_0} e^{-\Delta h_l} e^{-(l-1)\Delta H_0}$ with $H_l = H_0 - \sum_r \phi_{r,l} m_{z,r}$. Hence,

$$Z_b[\phi] = \det \left[1 + \prod_{l=0}^N e^{-\Delta h_l} \right] \equiv Z_f[\phi]. \quad (2)$$

We also demonstrate that the proposed formalism is equivalent to that of BSS [2]. Starting from the expression for Z found between Eq. (9) and Eq. (10) and using a matrix notation the partition function reads

$$\ln \frac{Z_b[\phi]}{Z_0} = \int_0^1 du \sum_{\sigma} \text{Tr} \Phi G_{\sigma}, \quad (3)$$

where Φ is a matrix with elements $\Phi(rl, r'l') = \phi_{r,l} \delta_{rr'} \delta_{ll'}$, the trace is over the space time indices and

$$G_{\sigma} = G^0 + u \sigma G^0 \Phi G_{\sigma} = (1 - u \sigma G^0 \Phi)^{-1} G^0, \quad (4)$$

where G^0 is the non interacting Green function. From Eqs. (3) and (4) we get

$$\ln \frac{Z_b[\phi]}{Z_0} = \int_0^1 du \sum_{\sigma} \text{Tr} \frac{1}{1 - u \sigma G^0 \Phi} \sigma G^0 \Phi. \quad (5)$$

The integral in terms of u and σ can be carried out analytically yielding the BSS partition function

$$Z_b[\phi] = Z_0 \exp(-\text{Tr}[\ln(1 - (G^0 \Phi)^2)]) \\ = \frac{Z_0}{\det(1 - (G^0 \Phi)^2)} = \prod_{\sigma=\uparrow\downarrow} \frac{1}{\det G_{\sigma}}. \quad (6)$$

A sign problem is present in Eq. (3), but is hidden in the u integral. If $G^0 \Phi$ has eigenvalues on the real axis with absolute value greater than 1, then the u integral runs over poles. This will give phases of $i\pi$ in the exponent which can lead to negative values of the exponential.

Finally, Efetov's *et al.* suggest an alternative way of evaluating the same weight Z_b from a bosonic field $A_{rr'}(\tau)$; however, the equation of motion that A satisfies, [their Eq. (13)], is singular. Furthermore, the equation of motion cannot uniquely determine A – not even with their additional constraint $\sum_r A_{rr}(\tau) = 0$. The latter can be shown, for example, using a two site cluster and a single time slice.

In conclusion, we have shown that for piecewise continuous paths the partition function obtained from Efetov *et al.*'s method [1] is equivalent to the standard BSS formulation and has, in particular, the same sign problem. However, their method provides a new perspective on the minus sign problem, as it can be viewed as originating from poles of the coupling constant (u) integral, or as branch cuts of a logarithm when the integral over u is performed analytically.

-
- [1] K. B. Efetov, C. Pépin, and H. Meier, Phys. Rev. Lett. **103**, 186403 (2009).
 [2] R. Blankenbecker, D. J. Scalapino, and R. L. Sugar, Phys. Rev. D **24**, 2278 (1981).